



B E C K E R
EAR, NOSE & THROAT CENTER

Princeton Office
256 Bunn Drive Suite A
(609) 430-9200

Robbinsville Office
1 Union Street Suite 203
(609) 436-5740

MEDICATION INSTRUCTIONS FOR ALLERGY PATIENTS

1. Drugs which contain antihistamine or have antihistaminic effects can result in negative reactions to skin testing. As a result, it may not be possible to properly interpret skin test results, and testing may have to be repeated at a later date. While this list is extensive, it is NOT all inclusive (particularly of the various brand names). Discontinue ALL antihistamines including the following medications **seven (7) days prior** to skin testing:

Antihistamines – Generic name (Brand name(s)):

Cetirizine (Zyrtec, Zyrtec-D)

Desloratidine (Clarinex)

Fexofenadine (Allegra, Allegra-D)

Hydroxyzine (Vistaril, Atarax)

Levocetirizine (Xyzal)

Loratidine (Claritin, Claritin-D)

Diphenhydramine (Aleve PM, Benadryl, Bayer P.M., Benylin, Contac P.M., Doans P.M., Excedrin PM, Legatrin P.M., Nytol, Tylenol Nighttime, Unisom, Zzzquil)

Chlorpheniramine (Aller-Chlor, Allerest, Alka Seltzer Plus, Chlor-Trimeton, Comtrex, Contac, Co-Pyronil, Coricidin, CTM, Deconamine, Dristan, Dura-tap, Naldecon, Ornade Spansules, Rondec, Sinutab, Teldrin, Triaminic, Triaminicin, Tylenol Allergy)

Azatadine (Optimine, Trinalin)

Brompheniramine (Bromfed, Dimetane, Dimetapp)

Carbinoxamine (Clistin, Rondec)

Clemastine (Tavist)

Cyclizine (Marezine)

Cyprohepatidine (Periactin)

Dexbrompheniramine (Drixoral)

Dexchlorpheniramine (Extendryl, Polaramine)

Dimenhydrinate (Dramamine)

Doxylamine (Nyquil)

Meclizine (Antivert)

Pheniramine

Phenyltoloxamine (Nadecon)

Promethazine (Phenergan)

Pyrilamine (Mepyramine)

Quinacrine (Atabrine)

Tripeleennamine (PBZ)

Tripolidine (Actidel, Actifed)

Antihistamines (Nasal Sprays)

Azelastine (Astelin, Astepro, Dymista)

Olopatadine (Patanase)

Antihistamines (Eye Drops)

Alcaftadine (Lastacft) Azelastine (Optivar) Bepotastine (Bepreve)
Emadastine (Emadine) Epinastine (Elestat) Ketotifen (Zaditor, Alaway)
Naphazoline/Pheniramine (Naphcon-A, Opcon-A, Visine-A)
Olopatadine (Patanol, Pataday, Pazeo)

Antihistamines (H2 blockers):

Cimetidine (Tagamet) Famotidine (Pepcid) Ranitidine (Zantac)

Tricyclic Antidepressants:

Amitriptyline (Elavil, Etrafon, Limbitrol, Triavil) Maprotiline (Ludiomil)
Amoxapine (Asenden) Mirtazapine (Remeron)
Clomipramine (Anafranil) Nortriptyline (Pamelor)
Desipramine (Norpramin) Protriptyline (Vivactil)
Doxepin (Silenor, Sinequan) Trimipramine (Surmontil)
Imipramine (Tofranil)

Do not take over-the counter cold medications that contain antihistamines. Read the label and look for “antihistamine” in the active ingredient list.

Drug Facts	Failure to follow these warnings could result in serious consequences.	Drug Facts (continued)	Drug Facts (continued)						
Active ingredients (in each 30 mL dose cup) Acetaminophen 650 mg Pain reliever/fever reducer Dextromethorphan HBr 30 mg Cough suppressant Doxylamine succinate 12.5 mg Antihistamine	Purpose temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> cough due to minor throat and bronchial irritation sore throat • headache minor aches and pains fever • runny nose and sneezing 	<ul style="list-style-type: none"> a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema trouble urinating due to enlarged prostate gland a sodium-restricted diet 	Directions <ul style="list-style-type: none"> take only as directed - see Overdose warning use dose cup or tablespoon (TBSP) do not exceed 4 doses per 24 hrs 						
Uses temporarily relieves common cold/flu symptoms: <ul style="list-style-type: none"> cough due to minor throat and bronchial irritation sore throat • headache minor aches and pains fever • runny nose and sneezing 		When using this product <ul style="list-style-type: none"> do not use more than directed excitability may occur, especially in children marked drowsiness may occur avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery 	<table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>30 mL (2 TBSP) every 6 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>	adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	30 mL (2 TBSP) every 6 hrs								
children 4 to under 12 yrs	ask a doctor								
children under 4 yrs	do not use								
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take			<ul style="list-style-type: none"> when using other DayQuil® or NyQuil products, carefully read each label to insure correct dosing 						

- You can continue all asthma medications, other eye drops, and other nasal sprays as prescribed. Also, continue all other medications for conditions other than allergies such as heart medications, blood pressure medications, etc.
- If your appointment date does not allow you to discontinue these medications 7 days in advance, please contact our front desk to reschedule (Princeton office phone is (609) 430-9200; Robbinsville office phone is (609) 436-5740).
- Remember to arrive 15 minutes early for paperwork and vital signs. Pediatric patients must be accompanied by a parent or medical guardian.